

A NOTE ON CHROMATICITY OF MULTIBRIDGE GRAPHS

A. M. KHALAF

Department of Mathematics, Faculty of Mathematics and Computer Science, University of Kufa, Najaf, Iraq

ABSTRACT

Coloring the vertices of graph such that every two adjacent vertices have different colors is a very common problem in graph theory. This is called a proper coloring of the graph. The possible number of different proper colorings on a graph with a given number of colors can be represented by a function called the chromatic polynomial. Two graphs G and H are said to be chromatically equivalent, if they share same chromatic polynomial. A graph G is chromatically unique, if G isomorphic H for any graph H such that G is chromatically equivalent H . The study of chromatically equivalent and chromatically unique problems called chromaticity. In this paper, we prove the chromatic uniqueness of a new family of multibridge graphs.

KEYWORDS: Graph & Chromatic Polynomial

Received: Jun 28, 2013; **Accepted:** Jul 20, 2013; **Published:** Aug 01, 2018; **Paper Id.:** IJMCARAUG20183

1. INTRODUCTION

The graphs that we consider here are finite, undirected and simple. Let $P(G, \lambda)$ denote the chromatic polynomial of a graph G . Two graphs G and H are said to be chromatically equivalent, and we write $G \sim H$, if $P(G, \lambda) = P(H, \lambda)$. A graph G is chromatically unique, unique if $G \cong H$ for any graph H such that $G \sim H$. By subdivision we mean the operation of replacing an edge of a graph by a path. If a graph H can be derived from G by a sequence of subdivisions, we say H is a subdivision of G . For each positive integer f , the graph $G(f)$ obtained from G by replacing each edge of G with a path of length f is called the f -uniform subdivision of G . The operation replaces a $u-v$ chain by an edge uv called chain-contraction. By contracting all maximal chains of a graph G , we arrive at multigraph $M(G)$. Two graphs G and H are homeomorphic if $M(G) = M(H)$. If G is homeomorphic to H we also say G is a H -homeomorph.

For each integer $k \geq 2$, let θ_k be the multigraph with two vertices and k edges. Any subdivision of θ_k is called multi-bridge graph or k -bridge graph. We denote $\theta(a_1, a_2, \dots, a_k)$ where $a_1, a_2, \dots, a_k \in \mathbb{N}$ to be the graph obtained by replacing the edges of θ_k by paths of length a_1, a_2, \dots, a_k respectively. Kalaf [6], Khalaf and Peng [7-11] showed the chromatic uniqueness of several kinds of k -bridge graphs.

We will solve a problem stated by Dong, Koh and Teo [4] related to chromaticity of 5-bridge graphs.

2. KNOWN RESULTS

In this section we cite some results previous results related to our objectives. A 2-bridge graph is simply cycle, which is χ -unique [4]. Chao and whithead Jr. [2], showed that every 3-bridge graph $\theta(1, a_2, a_3)$ called a theta graph is χ -unique. Loerinc [15] extended the above result to all 3-bridge graphs (called generalized θ graph). Bao and Chen [1] showed that every 5-bridge graph is χ -unique if its shortest maximal chains of length

greater than three. This result is special case of general result due to Xu, Liu and Peng [17].

Theorem 1 [17] For $k \geq 4$, $\theta(a_1, a_2, \dots, a_k)$ is χ – unique if $k - 1 \leq a_1 \leq a_2 \leq \dots \leq a_k$.

Li and Wei [14] established the following result:

Theorem 2 [14] For $a, b \in \mathbb{N}$, where $b \geq a \geq 3$, the 5-bridge graph $\theta(2, 2, 2, a, b)$ is χ – unique if and only if $(a, b) = (3, 4)$.

Ye [18] extended the above result to any k -bridge graph $\theta(2, 2, \dots, a, b)$ with $b \geq a \geq 3$ and $k \geq 5$. Li and Wei [14] stated without proof that $\theta(a, a, a, b, c)$, where $c \geq b \geq a + 1 \geq 3$, is χ – unique if and only if $(b, c) = (a + 1, a + 2)$. Xu, Liu, and Peng [17] showed that any f -uniform subdivision of θk is χ – unique, as in this theorem:

Theorem 3 [17] For $k \geq 2$, the graph $\theta k(f)$ is χ – unique.

The above result was proved independently by Dong [3], Koh and Teo [12] and Xu, Liu and Peng [17]. Peng [16] proved the following result:

Theorem 4 [16] Let $h, s \in \mathbb{N}$ such that $h \geq s + 1 \geq 2$ or $s = h + 1$. Then for $k \geq 4$, $\theta(a_1, a_2, \dots, a_k)$ is χ – unique if $a_2 - 1 = a_1 = h$, $a_j = h + s$ ($j = 3, \dots, k - 1$), $a_k \geq h + s$ and $a_k \in \{2h, 2h + s, 2h + s - 1\}$. Dong, Teo, Little, Hendy and Koh [5] proved this result:

Theorem 5 [5] If $2 \leq a_1 \leq a_2 \leq \dots \leq a_k < a_1 + a_2$, where $k \geq 3$, then the graph $\theta(a_1, a_2, \dots, a_k)$ is χ – unique.

Let $a_1, a_2, \dots, a_k \in \mathbb{N}$, and $G = \theta(a_1, a_2, \dots, a_k)$. Then see [4]:

$$P(G, \lambda) = \frac{1}{\lambda^{k-1}(\lambda-1)^{k-1}} \prod_{i=1}^k ((\lambda-1)^{a_i+1} + (-1)^{a_i+1}(\lambda-1)) + \frac{1}{\lambda^{k-1}} \prod_{i=1}^k ((\lambda-1)^{a_i} + (-1)^{a_i}(\lambda-1)).$$

Let $\lambda = 1 - x$, then:

$$\begin{aligned} P(G, 1-x) &= \frac{(-1)^{a_1+a_2+\dots+a_k+1}}{(1-x)^{k-1}} (x \prod_{i=1}^k (x^{a_i} - 1) - \prod_{i=1}^k (x^{a_i} - x)) \\ &= \frac{(-1)^{e(G)+1}}{(1-x)^{e(G)-v(G)+1}} (x \prod_{i=1}^k (x^{a_i} - 1) - \prod_{i=1}^k (x^{a_i} - x)), \end{aligned}$$

where $e(G) = \sum_{i=1}^k a_i$ and $v(G) = \sum_{i=1}^k a_i - k + 2$.

Also they defined $Q(G, x)$ for any graph G and real number x as:

$Q(G, x) = (-1)^{e(G)+1} (1-x)^{e(G)-v(G)+1} p(G, 1-x)$, and they got the following results:

Theorem 6 [4] for any $k, a_1, a_2, a_k \in \mathbb{N}$

$$Q(\theta(a_1, a_2, \dots, a_k, x)) = x \prod_{i=1}^k (x^{a_i} - 1) - \prod_{i=1}^k (x^{a_i} - x).$$

In particular, $Q(C_n, x) = x(x-1)(x^{n-1}-1)$.

Theorem 7 [4] for any two graphs G and H ,

(i) if $H \sim G$, then $Q(H, x) = Q(G, x)$;

(ii) if $Q(H, x) = Q(G, x)$ and $v(H) = v(G)$, then $H \sim G$.

Lemma 8 [4] suppose that $\theta(a_1, a_2, \dots, a_k) \sim \theta(b_1, b_2, \dots, b_k)$,

Where $k \geq 3, 2 \leq a_1 \leq a_2 \leq \dots \leq 2 \leq b_1 \leq b_2 \leq \dots \leq b_k$. then $a_i = b_i$ for all $i=1,2,\dots,k$.

Dong et al denote $g_e(G_1, G_2, \dots, G_k)$ to the collection of all edge-gluing of all G_1, G_2, \dots, G_k , wherer $k \geq 2$ and $e(G_i) \geq 1$ for all i , and then they got the following lemma:

Lemma 9 [4] Let $H \sim \theta(a_1, a_2, \dots, a_k)$, where $k \geq 3$ and $a_i \geq 2$ for all i .

Then one of the following is true:

$$1. H \cong \theta(a_1, a_2, \dots, a_k);$$

$$2. H \in g_e(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}} + 1, \dots, C_{b_{k+1}}), \text{ Where } 3 \leq t \leq k-1 \text{ and } b_i \geq 2 \text{ for all } i=1,2,\dots,k.$$

Theorem 10 [4] Let $k, t, b_1, b_2, \dots, b_k \in \mathbb{N}$ with $3 \leq t \leq k-1$ and $b_i \geq 2$ for all $i=1,2,\dots,k$.

If $H \in g_e(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}} + 1, \dots, C_{b_{k+1}})$ then

$$Q(H, x) = x \prod_{i=1}^k (x^{b_i} - 1) - \prod_{i=t+1}^t (x^{b_i} - x) \prod_{i=t+1}^k (x^{a_i} - 1).$$

Dong, Teo, Little, Hendy and Koh (2004) showed that the graph $\theta(2,2,2,3,4)$

Theorem 11 [5] $\theta(2,2,2,3,4) \sim H$ for every H the family $g_e \theta(2,2,2,3), C_4, C_4$.

3. RESULTS

In this section we solve a problem stated Dong, Koh and Teo [4] and give another result related to 6-bridge graph.

Problem 12 [4] Study the chromaticity of $\theta(a, a_2, \dots, a_5)$, where $2 \leq a_1 \leq a_2 \leq \dots \leq a_5 \leq 3$.

We completely solve the above problem by following proposition:

Proposition 13 The 5-bridge graph $\theta(a, a_2, \dots, a_5)$, where $2 \leq a_1 \leq a_2 \leq \dots \leq a_5 \leq 3$ is χ -unique.

Proof: There are six cases to be considered, depending on the length of a_i , $1 \leq i \leq 5$.

Case 1: The graph $\theta(a, a_2, \dots, a_5) = \theta(2,2,2,2,2)$, that is the graph $\theta_5(2)$ is x -unique by theorem 3.

Case 2: The graph $\theta(a, a_2, \dots, a_5) = \theta(2,2,2,2,3)$ is x -unique because this graph is one of x -unique graphs listed by Li [13].

Case 3: The graph $\theta(a, a_2, \dots, a_5) = \theta(2,2,2,3,3)$ is χ -unique because $(a, b) = (3,3) \neq (3,4)$ in the theorem 2.

Case 4: The graph $\theta(a, a_2, \dots, a_5) = \theta(2,2,3,3,3)$. In this graph $2 \leq a_1 \leq a_2 \leq \dots \leq a_5 \leq a_1 + a_2 = 4$.

By Theorem 5 is χ -unique.

Case 5: The graph $\theta(a_1, a_2, \dots, a_5) = \theta(2,3,3,3,3)$ choose $h=2$ and $s=1$, we see that $h \geq s+1=2$ and $a_1=2, a_2=3, a_3=3, a_4=3, a_5=3 \notin \{2h, 2h+s, 2h+s-1\}$. By Theorem 4, we conclude that $(2,3,3,3,3)$ is χ -unique.

Case 6: The graph $\theta(a, a_2, \dots, a_5) = \theta(3,3,3,3,3)$ that is graph $\theta_5(3)$ which is χ -unique By Theorem 3. ■

Lemma 14 Let $H \sim \theta(a_1, a_2, \dots, a_5)$, if $a_1 > 2, a_2 \leq a_2 \leq \dots \leq a_5$. Then one of the following is true:

$$(i) H \sim \theta(a_1, a_2, \dots, a_5);$$

(ii) $H \in g_e(\theta(b_1, b_2, b_3, b_4), C_{b_5} + 1)$, where $b_i \geq 2$ for all $i=1,2,\dots,5$.

Proof: Let $H \sim \theta(a_1, a_2, \dots, a_5)$ then by Lemma 2 of the following is true :

(i) $H \sim \theta(a_1, a_2, \dots, a_5)$;

(ii) $H \in g_e(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}} + 1, \dots, C_{b_{5+1}})$, where $3 \leq t \leq 4$ and $b_i \geq 2$ for all $i=1,2,\dots,5$

Now,

$$\begin{aligned} Q(G, x) &= x \prod_{i=1}^k (x^{a_i} - 1) - \prod_{i=1}^t (x^{a_i} - x) \\ &= -x^{a_1+1+a_2} + x^{1+a_2} - x^{a_3+4} - x^{4+a_4} + x^{3+a_4+a_5} - x^{a_5+1+a_2} - x^{a_3+2+a_1+a_2} - x^{a_3+a_4+a_5+a_1+a_2} - \\ &\quad x^{2+a_4+a_1+a_2} + x^{3+a_5+a_1} + x^{a_3+1} + x^{a_3+1} + x^{1+a_3+a_4+a_5+a_1+a_2} + x^{a_3+a_4+1+a_1} + x^{1+a_3+a_4+a_2} - x^{a_5+2+a_1+a_2} - \\ &\quad x^{2+a_4+a_5+a_2} + x^{a_3+a_4+1} + x^{a_3+1+a_5+a_1} + x^5 + x^{3+a_4+a_1} - x^{2+a_4+a_5+1} + x^{a_1+1} - x^{4+a_5} + x^{a_3+a_4+a_5+1} + \\ &\quad x^{a_3+a_4+3} - x^{a_3+a_4+a_5+2} + x^{1+a_3+a_5+a_2} - x^{1+a_4+a_5} + x^{a_3+3+a_2} - x^{a_3+a_4+2+a_2} - x^{a_3+2+a_5+a_2} - x^{a_3+a_4+2} + \\ &\quad x^{1+a_4+a_5+a_2} - x^{1+a_3+a_1} - x^{4+a_1} + x^{a_3+3+a_5} + x^{3+a_1+a_2} - x^{1+a_4+a_1} + x^{a_5+a_1+a_2} - \\ &\quad x^{1+a_3+a_2} + x^{3+a_5+a_2} + x^{a_3+3+a_1} + x^{1+a_3+a_1+a_2} + x^{3+a_4+a_2} - x^{a_3+2+a_5+a_1} - x^{4+a_2} + x^{1+a_4} + \\ &\quad x^{1+a_4+a_1+a_2} - x^{a_5+a_1+1} + x^{1+a_4+a_5+a_1} + x^{a_5+1} - x^{1+a_4+a_2} - x^{a_3+1+a_5} - x \end{aligned}$$

and $Q(H, x)$ depend on value of t . If $t=3$, then

$$\begin{aligned} Q(H, x) &= x \prod_{i=1}^5 (x^{b_i} - 1) - \prod_{i=1}^3 (b_i - x) \prod_{i=t+1}^k (x^{a_i} - 1) \\ &= x^{2+b_3+b_5} + x^{2+b_1+b_5} - x^{1+b_4+b_2} + x^{b_2+2+b_4} + x^{b_3+b_5+b_1+b_2} - x - x^{b_3+b_1+b_2} - x^{b_5+1+b_1} - x^{2+b_1} + \\ &\quad x^{2+b_3+b_4} + x^{b_4+1} + x^{1+b_4+b_5+b_1} - x^{2+b_3+b_4+b_5} - x^{2+b_4+b_5+b_1} - x^{3+b_5} + x^{1+b_3+b_4+b_5+b_1+b_2} + x^{2+b_4+b_1} + \\ &\quad x^{1+b_1} - x^{2+b_3} + x^{b_3+b_4+b_1+b_2} + x^{b_2+1} + x^{1+b_3+b_4+b_5} + x^{b_3+b_2+1+b_1} + x^{b_2+2+b_5} - x^{1+b_3+b_4+b_1+b_2} - \\ &\quad x^{1+b_3+b_5} - x^{1+b_3+b_4} + x^{b_5+1} - x^{3+b_4} + x^3 + x^{3+b_4+b_5} - x^{b_3+b_2+1+b_1} - x^{b_2+2} + x^{1+b_3} + x^{b_1+1+b_4+b_5} - \\ &\quad x^{b_4+b_5+1} - x^{b_2+2+b_4+b_5} - x^{b_3+b_4+b_5+b_1+b_2} - x^{1+b_4+b_1} - x^{1+b_5+b_2} \end{aligned}$$

If $t=4$, then

$$\begin{aligned} Q(H, x) &= x \prod_{i=1}^5 (x^{b_i} - 1) - \prod_{i=t+1}^3 (x^{b_5} - 1) \prod_{i=1}^4 (x^{b_i} - x) \\ &= -x^{1+b_4+b_2} + x^{b_2+2+b_4} - x^{3+b_3} - x - x^{b_5+1+b_1} + x^{2+b_3+b_4} + x^{b_4+1} + x^{1+b_4+b_5+b_1} - x^{2+b_3+b_4+b_5} - \\ &\quad x^{2+b_4+b_5+b_1} + x^{1+b_3+b_4+b_5+b_1+b_2} + x^{2+b_4+b_1} + x^{1+b_1} + x^{b_3+b_4+b_1+b_2} + x^{b_1+1} + x^{1+b_3+b_4+b_5} - \\ &\quad x^{1+b_3+b_4+b_1+b_2} - x^{1+b_3+b_5} - x^{1+b_3+b_4} + x^{b_5+1} - x^{3+b_4} + x^{3+b_4+b_5} + x^{1+b_3} + x^{b_2+1+b_4+b_5} - x^{b_4+b_5+1} - \\ &\quad x^{b_2+2+b_4+b_5} - x^{b_3+b_4+b_5+b_1+b_2} - x^{1+b_4+b_1} - x^{1+b_5+b_2} - x^{b_3+b_5+b_2+2} + x^4 + x^{1+b_5+b_1+b_2} - x^{4+b_5} - \\ &\quad x^{b_2+2+b_5+b_1} + x^{3+b_5+b_1} + x^{2+b_3+b_1} - x^{3+b_3+b_5} - x^{3+b_1} + x^{b_2+2+b_1} - x^{1+b_3+b_1} + x^{b_2+3+b_5} - x^{b_3+b_2+1} - \\ &\quad x^{b_2+3} + x^{b_3+b_2+2} - x^{1+b_1+b_2} + x^{1+b_3+b_5+b_1} + x^{b_3+b_2+1+b_5} \end{aligned}$$

It is easy to see that if $a > 2$ and $t=3$, then the term x^3 cannot be cancel with any other terms in $q(H, x)$ and $q(G, x)$.

Therefore either $H \cong \theta(a_1, a_2, \dots, a_5)$ or $H \in g_e(\theta(b_1, b_2, b_3, b_4), C_{b_5} + 1)$.

Lemma 15 if $a_1 > 3$, then $\theta(a_1, a_2, \dots, a_5)$ is χ -unique.

Proof Let $H \sim \theta(a_1, a_2, \dots, a_5)$. By lemma 3, either $H \cong \theta(a_1, a_2, \dots, a_5)$ or $H \in g_e(\theta(b_1, b_2, b_3, b_4), C_{b_5} + 1)$.

If $H \in g_e(\theta(b_1, b_2, b_3, b_4), C_{b_5} + 1)$, then

$$\begin{aligned} Q(H, x) = & -x^{1+b_4+b_2} + x^{b_2+2+b_4} - x^{3+b_3} - x - x^{b_5+1+b_1} + x^{2+b_3+b_4} + x^{b_4+1} + x^{1+b_4+b_5+b_1} - x^{2+b_3+b_4+b_5} - \\ & x^{2+b_4+b_5+b_1} + x^{1+b_3+b_4+b_5+b_1+b_2} + x^{2+b_4+b_1} + x^{1+b_1} + x^{b_3+b_4+b_1+b_2} + x^{b_1+1} + x^{1+b_3+b_4+b_5} - \\ & x^{1+b_3+b_4+b_1+b_2} - x^{1+b_3+b_5} - x^{1+b_3+b_4} + x^{b_5+1} - x^{3+b_4} + x^{3+b_4+b_5} + x^{1+b_3} + x^{b_2+1+b_4+b_5} - x^{b_4+b_5+1} - \\ & x^{b_2+2+b_4+b_5} - x^{b_3+b_4+b_5+b_1+b_2} - x^{1+b_4+b_1} - x^{1+b_5+b_2} - x^{b_3+b_5+b_2+2} + x^4 + x^{1+b_5+b_1+b_2} - x^{4+b_5} - \\ & x^{b_2+2+b_5+b_1} + x^{3+b_5+b_1} + x^{2+b_3+b_1} - x^{3+b_3+b_5} - x^{3+b_1} + x^{b_2+2+b_1} - x^{1+b_3+b_1} + x^{b_2+3+b_5} - x^{b_3+b_2+1} - \\ & x^{b_2+3} + x^{b_3+b_2+2} - x^{1+b_1+b_2} + x^{1+b_3+b_5+b_1} + x^{b_3+b_2+1+b_5} \end{aligned}$$

And $Q(G, x)$ as in Lemma 3.

The term x_4 in $Q(H, x)$ cannot be cancel with any other terms in $Q(H, x)$ or in $Q(G, x)$. Contradiction with Theorem 7.

In the same technique using in the above lemma, we prove the following result:

Theorem 16: The 6-bridge graph $\theta(a, a, a, a, b, c)$, $a \leq b \leq c$ is chromatically unique.

REFERENCES

1. X. W. Bao and X. E. Chen (1994). Chromaticity of the graph $\theta(a, b, c, d, e)$, (Chines, English and Chines summaries), J. Xinjing Univ. Natur. Sci. 11, 19-22.
2. C. Y. Chao and E. G. Whitehead Jr. (1979), Chromatic polynomial of a family of graphs. Ars Combin. 15, 111-129.
3. F. M. Dong, on chromatic uniqueness of two infinite families of graphs, J. graph Theory 17 (1993) 387-392.
4. F. M. Dong, K. M. Koh and K. L. Teo (2005), Chromatic polynomials and chromaticity of graphs, World Scientific publishing Co. Pte. Ltd., Singapore.
5. F. M. Dong, K. L. Teo, C. H. C. Little, M. D. Hendy and K. M. Koh (2004), chromatically unique multibridge graphs, Electronic J. of Combin. The ory 11, # R12.
6. A. M. Khalaf, "Chromaticity of certain k -bridge graphs", Ph. D. thesis, Universiti Putra Malaysia, 2010.
7. A. M. Khalaf and Y. H. Peng, Another Proof of Chromatically Unique 5-Bridge Graphs, International Journal of Applied Mathematics 22(6), 1009–1020 (2009).
8. A. M. Khalaf, Y. H. Peng and K. A. Atan, Chromaticity of 5-bridge Graphs (II), International Journal of Applied Mathematics 23, 791–808 (2010).
9. A. M. Khalaf and Y. H. Peng, The Chromatic Uniqueness of a Family of 6-Bridge Graphs, AKCE Journal of Graphs and Combinatorics 6(3), 393–400 (2009).
10. A. M. Khalaf and Y. H. Peng, Chromaticity of the 6-bridge Graph $\theta(a, a, a, b, c, c)$, International Journal of Applied Mathematics 22(7), 1087–1111 (2009).
11. A. M. Khalaf and Y. H. Peng, A Family of chromatically unique 6-bridge graphs, Ars Combinatoria 94, 211–220 (2010).
12. K. M. Koh and C. P. Teo, Some results on chromatically unique graphs, Proc. Asian Math. Conf. (World Scientific, Singapore, 1990) 258-262.
13. N. Z. Li, The list of chromatically unique graphs order seven and eight, Discrete Mathematics 172 (1997) 193-221.

14. Rani, H. A. *Optimization And Effectiveness Of Bridge Construction Development Based On Value Engineering*.
15. X. F. Li, and X. S. Wei (2001), *The chromatic uniqueness of a family of 5-bridge graphs (Chines)*, *J. Qinghai Normal Univ.*, no.2,12-17.
16. B. Loerinc (1978), *Chromatic uniqueness of the generalized θ –graphs*, *Dscrete Math.* 23, 313-316.
17. Y. H. Peng (1994), *On the coefficients of a graph and chromatic uniqueness of certain n-partition graphs*, in: *Combinatorics, Graph Theory, Algorithms and Applications (Beijing, 1993)*(World Scientific River Edge, NJ), 307-316.
18. S. J. Xu, J. J Liu and Y. H. Peng (1994), *The chromaticity of s-bridge graphs and related graphs*, *Discrete Math.*135,349-358.
19. C. F. Ye (2002), *The chromatic uniqueness of s-bridge graphs (Chines)*, *J. Xinjiang Univ. Natur. Sci.* 19, no. 3, 246-265.